

# Comparative Dose Efficacy Study of Atorvastatin Versus Simvastatin, Pravastatin, Lovastatin, and Fluvastatin in Patients With Hypercholesterolemia (The CURVES Study)

Peter Jones, MD, Stephanie Kafonek, MD, Irene Laurora, PharmD, and Donald Hunninghake, MD, for the CURVES Investigators\*

The objective of this multicenter, randomized, open-label, parallel-group, 8-week study was to evaluate the comparative dose efficacy of the 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase inhibitor atorvastatin 10, 20, 40, and 80 mg compared with simvastatin 10, 20, and 40 mg, pravastatin 10, 20, and 40 mg, lovastatin 20, 40, and 80 mg, and fluvastatin 20 and 40 mg. Investigators enrolled 534 hypercholesterolemic patients (low-density lipoprotein [LDL] cholesterol  $\geq 160$  mg/dl [4.2 mmol/L] and triglycerides  $\leq 400$  mg/dl [4.5 mmol/L]). The efficacy end points were mean percent change in plasma LDL cholesterol (primary), total cholesterol, triglycerides, and high-density lipoprotein cholesterol concentrations from baseline to the end of treatment (week 8). Atorvastatin 10, 20, and 40 mg

produced greater ( $p \leq 0.01$ ) reductions in LDL cholesterol,  $-38\%$ ,  $-46\%$ , and  $-51\%$ , respectively, than the milligram equivalent doses of simvastatin, pravastatin, lovastatin, and fluvastatin. Atorvastatin 10 mg produced LDL cholesterol reductions comparable to or greater than ( $p \leq 0.02$ ) simvastatin 10, 20, and 40 mg, pravastatin 10, 20, and 40 mg, lovastatin 20 and 40 mg, and fluvastatin 20 and 40 mg. Atorvastatin 10, 20, and 40 mg produced greater ( $p \leq 0.01$ ) reductions in total cholesterol than the milligram equivalent doses of simvastatin, pravastatin, lovastatin, and fluvastatin. All reductase inhibitors studied had similar tolerability. There were no incidences of persistent elevations in serum transaminases or myositis. ©1998 by Excerpta Medica, Inc. (Am J Cardiol 1998;81:582-587)

The Adult Treatment Panel of the National Cholesterol Education Program has established guidelines for the evaluation and treatment of elevated cholesterol concentrations based on an individual's risk factors for coronary artery disease.<sup>1</sup> The low-density lipoprotein (LDL) cholesterol treatment goals are (1) LDL cholesterol  $\leq 100$  mg/dl for patients with CAD; (2) LDL cholesterol  $< 130$  mg/dl in patients with  $\geq 2$  risk factors for CAD; (3) LDL cholesterol  $< 160$  mg/dl in patients with  $< 2$  risk factors for CAD. The Adult Treatment Panel recommended bile acid resins, nicotinic acid, and the 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase inhibitors as first-line drug treatments to achieve these treatment goals. Simvastatin, pravastatin, lovastatin, and fluvastatin lower LDL cholesterol from 18% to 41% over the most commonly used recommended dose range of each agent.<sup>2-13</sup> A recently approved synthetic HMG-CoA reductase inhibitor, atorvastatin, reduces LDL cholesterol from 35% to 61% over the dose range of

10 to 80 mg.<sup>14-18</sup> The present multicenter study (CURVES) was designed to evaluate the comparative dose efficacy of the HMG-CoA reductase inhibitor, atorvastatin, with equivalent dose strengths of simvastatin, pravastatin, lovastatin, and fluvastatin, in hypercholesterolemic patients after 8 weeks of treatment.

## METHODS

**Study design:** This study was a multicenter, open-label, randomized, parallel-group, 8-week comparative study evaluating the efficacy of once-daily doses of atorvastatin 10, 20, 40, and 80 mg compared with once-daily doses of simvastatin 10, 20, and 40 mg, pravastatin 10, 20, and 40 mg, lovastatin 20 and 40 mg, and fluvastatin 20 and 40 mg, and twice daily doses of lovastatin 40 mg (80 mg total daily dose). Male and female patients 18 to 80 years old with plasma LDL cholesterol concentrations  $\geq 160$  mg/dl (4.2 mmol/L) as calculated by the Friedewald formula, and triglyceride concentrations  $\leq 400$  mg/dl (4.5 mmol/L) at 2 consecutive visits (weeks -6 and -2) were eligible for inclusion.<sup>19</sup> Patients with any of the following conditions were excluded: primary hypothyroidism; nephrotic syndrome; type 1 or uncontrolled type 2 diabetes mellitus; hepatic dysfunction; serum creatine phosphokinase levels  $> 3$  times the upper limit of normal; body mass index  $> 32$  kg/m<sup>2</sup>; uncontrolled hypertension; myocardial infarction, coronary angioplasty, coronary artery bypass graft, or severe or unstable angina pectoris within the 3 months

From Baylor College of Medicine, The Methodist Hospital, Houston, Texas; Cardiovascular Medical Research, Parke-Davis, Division of Warner Lambert Company, Morris Plains, New Jersey; and Heart Disease Prevention Clinic, University of Minnesota School of Medicine, Minneapolis, Minnesota. This study was supported by Parke-Davis, Division of Warner Lambert Company, Morris Plains, New Jersey. Manuscript received August 20, 1997; revised manuscript received and accepted November 24, 1997.

Address for reprints: Peter H. Jones, MD, Baylor College of Medicine, 6565 Fannin #A601 Houston, Texas 77030.

\*See Appendix for list of CURVES investigators.

before the study; known hypersensitivities to HMG-CoA reductase inhibitors; or significant abnormalities that the investigator believed could compromise the patient's safety or successful participation in the study. Medications known to effect lipid levels, interact with study medications, or effect clinical laboratory parameters (erythromycin, anticoagulants, isotretinoin, immunosuppressive agents, lipid-regulating drugs, systemic steroids) were not allowed during the study.

Eligible patients were instructed to follow the step 1 diet for 6 weeks before randomization and throughout the duration of the study. After dietary stabilization, patients who qualified were randomized to 1 of 15 treatment groups, as described above, and were treated for 8 weeks. All study medication was taken according to recommended dosing. The study was performed using a common protocol at 34 sites. An appropriate institutional review board at all sites approved the protocol and all patients signed written informed consent.

**Laboratory methods:** Using standardized procedures, Medical Research Laboratories, Highland Heights, Kentucky, performed lipid and clinical laboratory measurements for all sites. The laboratory was certified for standardization of lipid analyses as specified by the Standardization Program of the Centers for Disease Control and Prevention and the National Heart, Lung, and Blood Institute.<sup>20</sup> After patients fasted overnight (minimum of 12 hours), blood was drawn in evacuated tubes containing ethylenediaminetetraacetic acid (1 mg/ml). Total plasma cholesterol and triglycerides were determined enzymatically with the Hitachi 747 analyzer (Boehringer Mannheim Diagnostics, Indianapolis, Indiana).<sup>21</sup> Plasma high-density lipoprotein (HDL) cholesterol was determined enzymatically after LDL and very low density lipoprotein cholesterol were selectively removed from the plasma by heparin and manganese chloride precipitation.<sup>22</sup> LDL cholesterol concentration was estimated by the Friedewald formula.<sup>19</sup> Fibrinogen was measured by immunonephelometry using an antiserum to human fibrinogen (BNA-100 Behring Diagnostics, Westwood, Massachusetts) in EDTA plasma stored at 70°C before analysis.

**Safety:** To monitor safety, complete clinical laboratory determinations were obtained at screening, randomization, and the end of the active treatment period. Physical examinations were performed at the beginning and end of the study. Adverse events were recorded at each clinic visit. Serum transaminases and creatinine phosphokinase concentrations were determined at every study visit and as deemed necessary by the investigator.

**Statistical methods:** Sample sizes were calculated based on the 2-sided Dunnett's test with a significance level of 5% and a standard deviation of 13% to detect differences in LDL cholesterol reductions of 8% (e.g., simvastatin 10 mg vs atorvastatin 10 mg) to 24% (e.g., fluvastatin 40 mg vs atorvastatin 40 mg) between atorvastatin and other reductase inhibitors at each dose level with at least 80% power.<sup>23</sup> The sample size in

each treatment arm varied greatly due to the large range of differences in lipid-lowering efficacy between atorvastatin and the other reductase inhibitors. Sample sizes were inflated by 5% for enrollment targets to allow for potential dropouts.

The intent-to-treat analysis performed for all efficacy end points included all randomized patients with post-treatment efficacy data for the primary efficacy end point of percent change in LDL cholesterol from baseline to week 8, and the secondary efficacy end points of percent change from baseline to week 8 in total cholesterol, triglycerides, and HDL cholesterol. Baseline was defined as the mean of measurements at week -2 and week 0 (randomization).

For each lipid parameter, the percent change was analyzed using an analysis of covariance model to test the treatment effect while controlling for baseline lipids. The least-squares means and mean square error from this model were used to compare atorvastatin with other reductase inhibitors at each dose level using Dunnett's procedure to fix the dose-wise type I error rate at 5%.<sup>23</sup> An analysis of variance model was used to test the assumption of no treatment-by-baseline lipid interaction.

Comparison between the least-squares means from the analysis of covariance model were used to evaluate (post hoc) the effect of each reductase inhibitor at all dose levels compared with atorvastatin 10 mg and atorvastatin 20 mg.

Safety was assessed among all patients receiving study medication using adverse events (coded using a modified COSTART dictionary) and clinical laboratory assessments. Particular attention focused on the presence of myopathy or elevated serum transaminase levels because these conditions have been associated with the use of reductase inhibitors.<sup>24</sup>

## RESULTS

**Patient characteristics:** Of the 534 patients randomized to treatment, 518 patients completed the study. Sixteen patients (3%) withdrew before the end of the study: 8 because of adverse events, 4 for personal reasons, and 4 who were lost to follow-up. The intent-to-treat analysis included 522 patients who provided post-treatment efficacy data. Fifty-nine percent of patients (307) were men and 41% (215) were women; 90% (469) were white. Mean age was 55 years (range 20 to 80), and 17% of patients had established CAD.


**Effects on serum lipids:** Mean baseline LDL cholesterol concentrations ranged from 192 to 244 mg/dl (5.0 to 6.3 mmol/L) and were similar across treatment groups (Table I). When given once daily in equivalent (mg) doses, atorvastatin 10, 20, and 40 mg produced greater ( $p \leq 0.01$ ) reductions in LDL cholesterol than simvastatin, pravastatin, lovastatin, and fluvastatin (Figure 1). Atorvastatin administered once daily at 80 mg reduced LDL cholesterol by 54%, whereas lovastatin administered as 40 mg twice daily reduced LDL cholesterol by 48%. This difference was not statistically significant ( $p = 0.17$ ) (Table II).

Atorvastatin 10 mg produced greater ( $p \leq 0.02$ ) reductions in LDL cholesterol than simvastatin 10 mg,

**TABLE I** Baseline Lipid and Lipoprotein Cholesterol Concentrations

| Treatment    | Dose (mg) | Number of Patients | Total Cholesterol | Triglycerides | HDL Cholesterol | LDL Cholesterol |
|--------------|-----------|--------------------|-------------------|---------------|-----------------|-----------------|
| Atorvastatin | 10        | 73                 | 298 (7.72)        | 169 (1.91)    | 51 (1.33)       | 213 (5.52)      |
| Pravastatin  | 10        | 14                 | 309 (8.00)        | 176 (1.98)    | 49 (1.26)       | 226 (5.83)      |
| Simvastatin  | 10        | 70                 | 289 (7.48)        | 157 (1.78)    | 51 (1.31)       | 207 (5.36)      |
| Atorvastatin | 20        | 51                 | 297 (7.68)        | 172 (1.94)    | 49 (1.28)       | 213 (5.51)      |
| Pravastatin  | 20        | 41                 | 315 (8.14)        | 147 (1.66)    | 48 (1.25)       | 237 (6.14)      |
| Simvastatin  | 20        | 49                 | 313 (8.09)        | 159 (1.79)    | 51 (1.32)       | 230 (5.95)      |
| Fluvastatin  | 20        | 12                 | 322 (8.34)        | 188 (2.12)    | 49 (1.26)       | 236 (6.10)      |
| Lovastatin   | 20        | 16                 | 334 (8.63)        | 192 (2.17)    | 51 (1.32)       | 244 (6.32)      |
| Atorvastatin | 40        | 61                 | 286 (7.40)        | 153 (1.73)    | 50 (1.29)       | 206 (5.32)      |
| Pravastatin  | 40        | 25                 | 299 (7.73)        | 172 (1.94)    | 49 (1.28)       | 215 (5.57)      |
| Simvastatin  | 40        | 61                 | 300 (7.77)        | 173 (1.95)    | 47 (1.20)       | 219 (5.66)      |
| Fluvastatin  | 40        | 12                 | 275 (7.12)        | 173 (1.95)    | 49 (1.26)       | 192 (4.97)      |
| Lovastatin   | 40        | 16                 | 301 (7.79)        | 172 (1.94)    | 49 (1.26)       | 219 (5.65)      |
| Atorvastatin | 80        | 10                 | 296 (7.65)        | 150 (1.69)    | 53 (1.37)       | 213 (5.51)      |
| Lovastatin   | 80        | 11                 | 306 (7.90)        | 200 (2.26)    | 47 (1.21)       | 219 (5.66)      |

Values are expressed as mean mg/dl (mmol/L).



**FIGURE 1.** Percent reduction in low-density lipoprotein cholesterol (LDL-C) after 8 weeks of treatment with atorvastatin (●), simvastatin (▲), pravastatin (◆), lovastatin (■), and fluvastatin (○). \* $p \leq 0.01$  versus atorvastatin at mg equivalent doses; † $p \leq 0.02$  versus atorvastatin 10 mg; ‡ $p \leq$  versus atorvastatin 20 mg.

pravastatin 10 and 20 mg, lovastatin 20 and 40 mg, and fluvastatin 20 and 40 mg (Table III). Atorvastatin 20 mg produced greater ( $p \leq 0.01$ ) reductions in LDL cholesterol than simvastatin 10, 20, and 40 mg, pravastatin 10, 20, and 40 mg, lovastatin 20 and 40 mg, and fluvastatin 20 and 40 mg (Table III).

As with LDL cholesterol, atorvastatin 10, 20, and 40 mg produced greater ( $p \leq 0.01$ ) reductions in total cholesterol than simvastatin, pravastatin, lovastatin, and fluvastatin at milligram-equivalent doses (Table II). The effects on triglycerides were not different between atorvastatin and the other reductase inhibitors except at the 40-mg dose when atorvastatin produced greater ( $p \leq 0.05$ ) reductions in triglycerides than the 40-mg doses of simvastatin, pravastatin, lovastatin, and fluvastatin (Table II). Effects on HDL cholesterol, ranging from 3.0% to 9.9%, were not different between atorvastatin and the other reductase inhibitors except at the 40-mg

dose when simvastatin produced greater ( $p \leq 0.05$ ) elevations in HDL cholesterol than atorvastatin (Table II).

**Safety:** The overall frequency of adverse events was similar between treatment groups. Fifty-two patients (10%) reported adverse events that were judged by the investigator to be possibly, probably, or definitely associated with treatment, most of which were mild to moderate in intensity. Of these, the most commonly reported events were myalgia (1.5%), abdominal pain (1.3%), diarrhea (1.1%), flatulence (1%), and nausea (1%). Eight patients withdrew from the study due to adverse events: 2 in the atorvastatin group (1%), 4 in the simvastatin group (2%), and 1 each in the pravastatin (1%) and fluvastatin groups (4%) (Table IV). The adverse

events leading to withdrawal included gastrointestinal complaints, dizziness, depression, myalgia, hypertension, angina, and back pain.

There were no incidences of persistent (2 measurements within 1 week) elevations in serum transaminases  $>3$  times the upper limit of normal. There were no incidences of elevations in creatine phosphokinase  $>3$  times the upper limit of normal or reports of myopathy in any treatment group. There were no significant changes from baseline in mean fibrinogen levels for any of the reductase inhibitors.

## DISCUSSION

The CURVES study is the first trial to compare the lipid-lowering efficacy of all marketed HMG-CoA reductase inhibitors, including the recently approved synthetic HMG-CoA reductase, atorvastatin, across their dose ranges. An open-label design was chosen

**TABLE II** Mean Percent Change ( $\pm$ SD) in Lipoprotein Concentrations

| Treatment    | Dose (mg) | Number of Patients | Total Cholesterol     | Triglycerides         | HDL Cholesterol | LDL Cholesterol         |
|--------------|-----------|--------------------|-----------------------|-----------------------|-----------------|-------------------------|
| Atorvastatin | 10        | 73                 | -28 (9)               | -13 (25)              | 5.5 (12)        | -38 (10)                |
| Pravastatin  | 10        | 14                 | -13 (12) <sup>†</sup> | 3 (46)                | 9.9 (13)        | -19 (14) <sup>†</sup>   |
| Simvastatin  | 10        | 70                 | -21 (9) <sup>†</sup>  | -12 (30)              | 6.8 (9)         | -28 (12) <sup>†</sup>   |
| Atorvastatin | 20        | 51                 | -35 (6)               | -20 (25)              | 5.1 (11)        | -46 (8)                 |
| Pravastatin  | 20        | 41                 | -18 (7) <sup>†</sup>  | -15 (17)              | 3.0 (8)         | -24 (9)** <sup>†</sup>  |
| Simvastatin  | 20        | 49                 | -26 (8) <sup>†</sup>  | -17 (22)              | 5.2 (10)        | -35 (11)**              |
| Fluvastatin  | 20        | 12                 | -13 (6) <sup>†</sup>  | -5 (32)               | 0.9 (8)         | -17 (8)** <sup>†</sup>  |
| Lovastatin   | 20        | 16                 | -21 (9) <sup>†</sup>  | -12 (23)              | 7.3 (12)        | -29 (13)** <sup>†</sup> |
| Atorvastatin | 40        | 61                 | -40 (8)               | -32 (19)              | 4.8 (12)        | -51 (10)                |
| Pravastatin  | 40        | 25                 | -24 (7) <sup>†</sup>  | -10 (22) <sup>†</sup> | 6.2 (11)        | -34 (9)** <sup>†</sup>  |
| Simvastatin  | 40        | 61                 | -30 (10) <sup>†</sup> | -15 (29) <sup>†</sup> | 9.6 (13)*       | -41 (13)** <sup>†</sup> |
| Fluvastatin  | 40        | 12                 | -19 (9) <sup>†</sup>  | -13 (34)*             | -3.0 (10)       | -23 (10)** <sup>†</sup> |
| Lovastatin   | 40        | 16                 | -23 (6) <sup>†</sup>  | -2 (27) <sup>†</sup>  | 4.6 (13)        | -31 (7)** <sup>†</sup>  |
| Atorvastatin | 80        | 10                 | -42 (7)               | -25 (22)              | -0.1 (9)        | -54 (9)                 |
| Lovastatin   | 80        | 11                 | -36 (6)               | -13 (28)              | 8.0 (13)        | -48 (8)                 |

\* $p \leq 0.05$ ; <sup>†</sup> $p \leq 0.01$ , Dunnett's test of significance compared with atorvastatin at milligram-equivalent doses.

<sup>†</sup>Atorvastatin 10 mg statistically significantly better ( $p \leq 0.02$ ).

<sup>†</sup>Atorvastatin 20 mg statistically significantly better ( $p \leq 0.01$ ).

Values are expressed as mean percent change from baseline.

**TABLE III** Comparison of Percent Change in Low-Density Lipoprotein (LDL) Cholesterol: Atorvastatin 10 and 20 mg Versus All Treatments

| Treatment Group | Dose (mg) | Number of Patients | Mean* Percent Change from Baseline LDL Cholesterol | p Value vs Atorvastatin 10 mg | p Value vs Atorvastatin 20 mg |
|-----------------|-----------|--------------------|--|-------------------------------|-------------------------------|
| Atorvastatin    | 10        | 73                 | -38  | Referent                      | —                             |
| Atorvastatin    | 20        | 51                 | -46  | —                             | Referent                      |
| Fluvastatin     | 20        | 12                 | -17  | 0.0001                        | 0.0001                        |
| Fluvastatin     | 40        | 12                 | -23  | 0.0001                        | 0.0001                        |
| Lovastatin      | 20        | 16                 | -29  | 0.0019                        | 0.0001                        |
| Lovastatin      | 40        | 16                 | -31  | 0.0197                        | 0.0001                        |
| Lovastatin      | 80        | 11                 | -48  | NS                            | NS                            |
| Pravastatin     | 10        | 14                 | -19  | 0.0001                        | 0.0001                        |
| Pravastatin     | 20        | 41                 | -24  | 0.0001                        | 0.0001                        |
| Pravastatin     | 40        | 25                 | -34  | NS                            | 0.0001                        |
| Simvastatin     | 10        | 70                 | -28  | 0.0001                        | 0.0001                        |
| Simvastatin     | 20        | 49                 | -35  | NS                            | 0.0001                        |
| Simvastatin     | 40        | 61                 | -41  | NS                            | 0.0083                        |

\*Least-squares mean.

NS = atorvastatin not statistically significantly better.

for this study because of the impracticality of blinding 15 treatment arms. Efficacy end points were based on objective laboratory measurements.

Atorvastatin 10, 20, and 40 mg produced greater ( $p \leq 0.01$ ) reductions in total and LDL cholesterol than the other reductase inhibitors studied at milligram-equivalent doses. Atorvastatin 10 mg produced greater ( $p \leq 0.02$ ) reductions in LDL cholesterol than to simvastatin 10 mg, pravastatin 10 and 20 mg, lovastatin 20 and 40 mg, and fluvastatin 20 and 40 mg. The reduction in LDL cholesterol with atorvastatin 80 mg once daily ( $-54\%$ ) was numerically, but not statistically, greater than lovastatin administered as 40 mg twice daily ( $-48\%$ ) in a small sample of 10 and 11 patients, respectively.

The lipid-lowering effects observed in the present study are consistent with those seen in previous comparisons between HMG-CoA reductase inhibitors. Simvastatin 10 to 40 mg produced reductions in LDL cholesterol of 28% to 41%, pravastatin 10 to 40 mg

produced reductions in LDL cholesterol of 18% to 34%, lovastatin 20 to 40 mg produced reductions in LDL cholesterol of 25% to 38%, and fluvastatin 20 to 40 mg produced reductions in LDL cholesterol of 18% to 27%.<sup>2-13</sup> Only the lovastatin 40 mg twice-a-day treatment group had a greater reduction in LDL cholesterol in this study (48%) than anticipated based on the results from a large clinical trial—Expanded Clinical Evaluation of Lovastatin (EXCEL)—in which reductions were reported as 40%.<sup>25</sup> The greater than expected LDL cholesterol reductions in this group may be partially explained by the small sample size.

An HMG-CoA reductase inhibitor's efficacy is measured by its ability to lower LDL cholesterol regardless of the amount of drug substance needed to accomplish this result (potency). Atorvastatin, administered in doses of 10 to 80 mg to patients with primary hypercholesterolemia, lowers LDL cholesterol by 35% to 61%.<sup>14-18</sup> The present study, in conjunction with previous comparative studies that have

**TABLE IV** Withdrawals Due to Adverse Events

| Treatment    | Dose (mg) | No. of Patients | No. of Patients Withdrawn Due to Adverse Events | Event(s)                                       | Relation to Therapy* |
|--------------|-----------|-----------------|---|--|----------------------|
| Atorvastatin | 10        | 74              | 1   | Abdominal pain/diarrhea                        | Possibly             |
| Pravastatin  | 10        | 14              | 0   |  |                      |
| Simvastatin  | 10        | 70              | 1   | Depression/dizziness                           | Possibly             |
| Atorvastatin | 20        | 51              | 1   | Myalgia  | Definitely not       |
| Pravastatin  | 20        | 42              | 1   | Dizziness                                      | Probably             |
| Simvastatin  | 20        | 51              | 2   | Hypertonia/nausea<br>Abdominal pain/flatulence | Possibly<br>Probably |
| Fluvastatin  | 20        | 12              | 0   |  |                      |
| Lovastatin   | 20        | 16              | 0   |  |                      |
| Atorvastatin | 40        | 61              | 0   |  |                      |
| Pravastatin  | 40        | 25              | 0   |  |                      |
| Simvastatin  | 40        | 61              | 1   | Angina   | Unlikely             |
| Fluvastatin  | 40        | 12              | 1   | Back pain                                      | Probably             |
| Lovastatin   | 40        | 16              | 0   |  |                      |
| Atorvastatin | 80        | 10              | 0   |  |                      |
| Lovastatin   | 80        | 11              | 0   |  |                      |

\*Relation to therapy was judged by the investigator.

included atorvastatin, have clearly established atorvastatin as the most efficacious HMG-CoA reductase inhibitor for lowering LDL cholesterol.<sup>16-18</sup>

This study was not powered to detect differences in effects on triglycerides. The patient population studied consisted mostly (74%) of patients with elevated cholesterol without elevated triglycerides (mean baseline triglycerides ranged from 147 to 200 mg/dl [1.66 to 2.26 mmol/L]). Atorvastatin 10, 20, and 80 mg produced numerically, but not statistically, greater reductions in triglycerides than the other reductase inhibitors at milligram-equivalent doses, and statistically greater reductions in triglycerides at the 40 mg dose. As with LDL cholesterol, the reductions in triglycerides seen in all of the treatment groups in the present study are consistent with those reported in previous studies.<sup>2-18</sup>

Reductase inhibitors are generally well tolerated.<sup>24</sup> Clinically important adverse effects of the drugs include increases in serum transaminase concentrations and myositis, with or without complicating rhabdomyolysis. In the present study, no patient in any treatment arm experienced persistent clinically significant increases in serum transaminases. Most cases of significant elevations in serum transaminases have been reported to occur within the first 2 to 5 months of treatment, and the duration of this study (8 weeks) may not have been long enough to detect such cases.<sup>26</sup> In rare instances, severe creatine phosphokinase elevations (>10 times the upper limit of normal) and myositis have been associated with the use of reductase inhibitors.<sup>27</sup> In the present study, no subject experienced creatine phosphokinase concentrations >3 times the upper limit of normal, or myopathy.

**Acknowledgment:** We acknowledge and thank Boston Biostatistics Inc., Framingham, Massachusetts, for

their support and efforts related to the conduct of the study and the data analysis.

## APPENDIX

**CURVES Investigators:** W. Virgil Brown, MD, Emory University School of Medicine, Atlanta, GA; Arthur Bucci, MD, Mercy Heart Institute, Pittsburgh, PA; David Capuzzi, MD, PhD, Medical College of Pennsylvania, Philadelphia, PA; Albert Carr, MD, Southeastern Clinical Research & Management, Inc., Augusta, GA; Michael Clearfield, DO, University of North Texas, Fort Worth, TX; Stephen Crespino, MD, St. Louis, MO; Paresch Dandona, MD, State University of New York at Buffalo, Buffalo, NY; Michael Davidson, MD, Chicago Center for Clinical Research Inc., Chicago, IL; Fred Faas, MD, John L. McClellan Memorial Veterans Hospital, Little Rock, AR; Keith Ferdinand, MD, Margo Morgan Research Center, New Orleans, LA; Geoffrey S. Ginsburg, MD, PhD, Beth Israel Hospital, Boston, MA; Donald B. Hunninghake, MD, University of Minnesota, Minneapolis, MN; William Insull, MD, Baylor College of Medicine, The Methodist Hospital, Houston, TX; Peter H. Jones, MD, Baylor College of Medicine, The Methodist Hospital, Houston, TX; Stephanie Kafonek,\* MD, The Johns Hopkins University, Baltimore, MD; John P. Kane, MD, University of California, San Francisco, CA; Moti L. Kashyap, MD, Veterans Administration Medical Center, Long Beach, CA; Kent D. Katz, MD, Veterans Administration Medical Center, Long Beach, CA; Robert H. Knopp, MD, University of Washington, Harborview Medical Center, Seattle, WA; Peter Kwiterovich, MD, The Johns Hopkins University, Baltimore, MD; Andrew J. Lewin, MD, National Research Institute, Los Angeles, CA; Irving K. Loh, MD, Ventura Heart Institute, Thousand Oaks, CA; Charles P. Lucas, MD, William Beaumont Hospital, Birmingham, MI; James M. McKenney, PharmD, National Clinical Research Inc., Richmond, VA; John M. Morgan, Medical College of Pennsylvania, Philadelphia, PA; David T. Nash, MD, Syracuse, NY; Stephen D. Nash, MD, Syracuse, NY; Christopher M. Rembold, MD, University of Virginia, Charlottesville, VA; Lawrence M. Resnick, MD, Veterans Affairs Medical Center, Allen Park, MI; Robert G. Robertson, MD, Emory University School of Medicine, Atlanta, GA; Robert J. Rosenson, MD, Rush-Presbyterian/St. Luke's Medical Center, Chicago, IL; F. Julie Samuels, MD, National Clinical Research Inc., Richmond, VA; Xavier Pi-Sunyer, MD, St. Luke's/Roosevelt Hospital Center, New York, NY; Arkady Synhavsky, MD, Kidney Disease & Critical Care, Roseville, MN; Stephen F. Weis, DO, University of North Texas, Fort Worth, TX; Stuart R. Weiss, MD, The San Diego Endocrine & Medical Clinic Inc., San Diego, CA; James H. Zavoral MD, Preventive Cardiology Institute, Fairview Southdale Hospital, Edina, MN; Paul Ziajka, MD, PhD, The Florida Lipid Associates, Orlando, FL; Jean Bergeron, MD, Hotel-Dieu de Quebec, Quebec, Canada; Jacques Genest, MD, Clinical Research Institute of Montreal, Montreal, Quebec, Canada; Ruth McPherson, MD, PhD, University of Ottawa Heart Institute, Ottawa, Ontario Canada.

1. Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Summary of the Second Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel II). *JAMA* 1993;269:3015-3023.
2. Jones PH, Farmer JA, Cressman MD, McKenney JM, Wright JT, Proctor JD, Berkson DM, Farnham DJ, Wolfson PM, Colfer HT, Rackley CE, Sigmund WR, Schlant RC, Arenberg D, McGovern ME. Once-daily pravastatin in patients with primary hypercholesterolemia: a dose response study. *Clin Cardiol* 1991;14:146-151.
3. Illingworth DR. HMG CoA reductase inhibitors. *Curr Opin Lipidol* 1991;2:24-30.
4. Illingworth DR, Tobert JA. A review of clinical trials comparing HMG CoA reductase inhibitors. *Clin Ther* 1994;16:366-385.
5. Weir MR, Berger ML, Weeks ML, Liss CL, Santanello NC, for the Quality of Life Multicenter Group. Comparison of the effects on quality of life and the efficacy and tolerability of lovastatin versus pravastatin. *Am J Cardiol* 1996;77:475-479.
6. The Simvastatin Pravastatin Study Group. Comparison of the efficacy, safety and tolerability of simvastatin and pravastatin for hypercholesterolemia. *Am J Cardiol* 1993;71:1408-1414.
7. Simvastatin Pravastatin European Study Group. Comparative efficacy and

\*Stephanie Kafonek, MD, is now Senior Director of Cardiovascular Medical Research at Parke-Davis, a Division of Warner-Lambert.

- tolerability of 5 and 10 mg simvastatin and 10 mg pravastatin in moderate primary hypercholesterolemia. *Cardiology* 1994;85:244–254.
8. Douste-Blazy P, Ribeiro VG, Seed M, and the European Study Group. Comparative study of the efficacy and tolerability of simvastatin and pravastatin in patients with primary hypercholesterolemia. *Drug Invest* 1993;6:353–361.
  9. Illingworth DR, Stein EA, Knopp RH, Hunnighake DB, Davidson MH, Dujovne CA, Miller VT, Tobert JA, Laskarzewski PM, Isaacsohn JL, Bacon SP, Tate AC. A randomized multicenter trial comparing the efficacy of simvastatin and fluvastatin. *J Cardiovasc Pharmacol Ther* 1996;1:23–30.
  10. Berger ML, Wilson HM, Liss CL. A comparison of the tolerability and efficacy of lovastatin 20 mg and fluvastatin 20 mg in the treatment of primary hypercholesterolemia. *J Cardiovasc Pharmacol Therapeut* 1996;1:101–106.
  11. Jacotot B, Benghozi R, Pfister P, Holmes D, on behalf of the french fluvastatin study group. Comparison of fluvastatin versus pravastatin treatment of primary hypercholesterolemia. *Am J Cardiol* 1995;76:54A–56A.
  12. Schulte K, Beil S. Efficacy and tolerability of fluvastatin and simvastatin in hypercholesterolemic patients. *Clin Drug Invest* 1996;12:119–126.
  13. Lintott CJ, Sutherland WHF, Scott RS, Bremer J. Treating hypercholesterolemia with HMG CoA reductase inhibitors: a direct comparison of simvastatin and pravastatin. *Aust N Z J Med* 1993;23:381–386.
  14. Nawrocki JW, Weiss SR, Davidson MH, Sprecher DL, Schwartz SL, Lupien PJ, Jones PH, Haber HE, Black DM. Reduction of LDL-cholesterol by 25% to 60% in patients with primary hypercholesterolemia by atorvastatin, a new HMG-CoA reductase inhibitor. *Arterioscler Thromb Vasc Biol* 1995;15:678–682.
  15. Gmerek A, Yang R, Bays H, Jones P, Knopp R, Littlejohn T, Schrott H, Black D. Atorvastatin causes a dose-dependent reduction in LDL-C and triglycerides. In: 66<sup>th</sup> Congress of the European Artherosclerosis Society, July 1996:212.
  16. Davidson M, McKinney J, Stein E, Schrott H, Bakker-Arkema R, Fayyad R, Black D, for the Atorvastatin Study Group I. Comparison of the one-year efficacy and safety of atorvastatin versus lovastatin in primary hypercholesterolemia. *Am J Cardiol* 1997;79:1475.
  17. Bertolini S, Bittolo Bon G, Cambell LM, Farnier M, Langan J, Mahla G, Pauciullo P, Sirtori C, Egros F, Fayyad R, Nawrocki JW. Efficacy and safety of atorvastatin compared to pravastatin in patients with hypercholesterolemia. *Atherosclerosis* 1997;130:191–197.
  18. Dart A, Jerums G, Nicholson G, d'Emden M, Hamilton-Craig I, Tallis G, Best J, West M, Sullivan D, Bracs P, Black D. A multicenter, double-blind, one-year study comparing safety and efficacy of atorvastatin versus simvastatin in patients with hypercholesterolemia. *Am J Cardiol* 1997;80:39–44.
  19. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 1972;18:499–502.
  20. Myers GL, Copper GR, Winn CL, Smith SJ. The Centers for Disease Control—National Heart, Lung, and Blood Institute Lipid Standardization Program: an approach to accurate and precise lipid measurements. *Clin Lab Med* 1989;9:105–135.
  21. Steiner PM, Freidel J, Bremmer WF, Stein EA. Standardization of micromethods for plasma cholesterol, triglyceride and HDL-cholesterol with the Lipid Research Clinics' methodology (abstr). *J Clin Chem Clin Biochem* 1981;19:850.
  22. Steele WB, Koehle DF, Azar MM, Blaszkowski TP, Kuba K, Dempsey ME. Enzymatic determinations of cholesterol in high density lipoprotein fractions prepared by a precipitation technique. *Clin Chem* 1976;22:98–101.
  23. Dunnett CW. A multiple comparison procedure for comparing several treatments with a control. *J Am Stat Assoc* 1955;50:1096–1121.
  24. Blum C. Comparison of properties of four inhibitors of 3-hydroxy-3-methylglutaryl-coenzyme A reductase. *Am J Cardiol* 1994;73:3D–11D.
  25. Bradford RH, Shear CL, Chremos AN, Dujovne C, Downton M, Franklin FA, Gould AL, Hesney M, Higgins J, Hurley DP, Langendorfer A, Nash DT, Pool JL, Schnaper H. Expanded Clinical Evaluation of Lovastatin (EXCEL) study results. I. Efficacy in modifying plasma lipoproteins and adverse events profile in 8245 patients with moderate hypercholesterolemia. *Arch Intern Med* 1991;151:43–49.
  26. Hsu I, Spinler SA, Johnson NE. Comparative evaluation of the safety and efficacy of HMG-CoA reductase inhibitor monotherapy in the treatment of primary hypercholesterolemia. *Ann Pharmacother* 1995;29:743–759.
  27. Dujovne CA, Chremos N, Pool JL, Schnaper H, Bradfrd RH, Shear CL, Higgins J, Downton M, Franklin FA, Nash DT, Gould L, Langendorfer A. Expanded Clinical Evaluation of Lovastatin (EXCEL) study. IV. Additional perspectives on the tolerability of lovastatin. *Am J Med* 1991;91(suppl B):25S–30S.